

## ***Physa mezzalirai*, a new cretaceous basommatophoran from Adamantina formation, Brazil**

Renato P. Ghilardi<sup>1</sup>, Fábio A. Carbonaro<sup>1</sup> & Luiz Ricardo L. Simone<sup>2</sup>

<sup>1</sup>Laboratório de Paleontologia de Macroinvertebrados (LAPALMA), Universidade Estadual Paulista Júlio de Mesquita Filho.  
Av. Eng. Luiz Edmundo Carrijo Coube, 14-01, 17033-360, Bauru, SP, Brasil. ghilardi@fc.unesp.br, fabiocarbonaro@yahoo.com.br  
<sup>2</sup>Museu de Zoologia da Universidade de São Paulo, Cx. Postal 42494, 04218-970, São Paulo, SP, Brasil. lrsimone@usp.br

### **Abstract**

This paper deals with a formal description of *Physa mezzalirai* (Basommatophora, Physidae), a new species of freshwater gastropod that lived in upper Cretaceous deposits, in sand-silty beds of Adamantina formation, Bauru basin, from Marília, São Paulo state, Brazil. This formal description legitimates the availability of the name previously described in 2010 by same authors; being this paper the correct citation for the species name.

Keywords: *Physa mezzalirai* new species, Cretaceous, Brazil

### **Resumo**

Este artigo relata a descrição formal de *Physa mezzalirai* (Basommatophora, Physidae), uma espécie nova de gastrópode de água doce que viveu no cretáceo superior, depósitos arenoso-sílticos da formação Adamantina, Grupo Bauru, de Marília, São Paulo, Brasil. Esta descrição formal legitima a validade do nome previamente descrito em 2010 pelos mesmos autores, sendo este artigo a citação correta para o nome da espécie.

Palavras-chave: *Physa mezzalirai* nova espécie, Cretáceo, Brasil.

### **Introduction**

*Physa mezzalirai* (Basommatophora, Physidae), was already described by a previous paper, Ghilardi, Carbonaro & Simone (2010). However, that paper does not attend the ICZN article 8.6, causing the necessity of this new paper on the same taxon. This paper is, then, a formal, ruled description, which must be mentioned as actual reference of the species' name. A more complete description and an abstract on the geology of the Adamantina formation, and of the Bauru basin, can be found in Ghilardi et al. (2010: 63-65, figs. 1-2). Bauru group is rich in fossil invertebrates, mainly freshwater bivalves. The finding of gastropods is, on the other hand, rare, both, in species and specimens. The genus *Physa* Draparnaud, 1801 is practically the single of the family Physidae (Pulmonata, Basommatophora), it occurs worldwide in tropical and subtropical freshwater environments, and it is characterized by sinistral (left-hand coiled), small snails. Representatives occur since the Cretaceous to Recent (Taylor, 2003). Another congener species occur in Bauru group: *Physa aridi* Mezzalira, 1974.

### **Systematics**

**Order: Basommatophora**  
Family Physidae


***Physa mezzalirai*, new species**  
(Figs. 1-8)

*Physa mezzalirai* Ghilardi, Carbonaro & Simone, 2010: 63-67 (figs. 4A-H).

**Types:** Holotype MPM075 (Museu Municipal de Paleontologia de Marília). Paratype MPM076, same locality (copies in MZSP – Museu de Zoologia da Universidade de São Paulo).

**Type locality:** BRAZIL. São Paulo; outcrops in road Riacho Água – Formosa, 10 km south from Marília town, 22°20'28.14"S 49°56'41.94"W.

**Stratigraphy:** Adamantina formation, Bauru group, Turonian-Santonian, upper Cretaceous.



Figs 1-8 Types of *Physa mezzalirai* n. sp.. MPM075 holotype, 1, left view; 2, apertural view; 3, dorsal view; 4, apical view; 5, counter-mould kept in matrix; 6, detail of apical region in profile; 7, paratype, inner mould with remaining of shell in region of outer lip (center-inferior in Fig.); 8, detail of region between penultimate and last whorl, showing micro-sculpture close to suture.

## Description

Shell from 11 to 14 mm long, elliptical-elongated, adolute. Width ~47% of length (Figs. 1-3). Spire with 6 whorls, apex pointed at ~30° (Figs. 5, 6). Body whorl wider in middle level, occupying ~7/10 from total length (Figs. 2, 5); dorso-ventral height ~70% of width (Fig. 4). Suture shallow, almost planar. Growth lines weakly visible; sculptured by lyrate, axial, delicate, shallow ribs (Fig. 8), clearer in region between spire and body whorl; this lyrate sculpture fainting from middle body whorl to inferior region of it. Aperture orthocline, about twice longer than wide, occupying ~70% of length (Figs. 1, 2, 5, 7). Umbilicus wanting.

**Measurements (in mm):** Holotype (Figs. 1-6, 8): 11.2 by 8.0; paratype (Fig. 7): 14.0 by 6.5.

**Etymology:** the species epithet refers to Dr. Sergio Mezzalira, a great contributor to the knowledge on Bauru group fossils.

## Discussion

*Physa mezzalirai* differs from fossil species *P. wickmanni* Parodiz, 1961 and *P. doeringi* Doello-Jurado, 1927, both from Argentine Cretaceous, and from *P. aridi*, from São José do Rio Preto region, in having the wider proportion length/width of 2.24, while the remaining species are, respectively, 1.9, 1.3-1.7, and 1.7. The body whorl of *P. mezzalirai* is ~72%, while the other species are, respectively, 74, 79 and 75%. Related to the number of whorls, *P. mezzalirai* resembles *P. wickmanni*, in having ~6; but it differs from the other two species, which have 4-5 whorls (for additional morphometric parameters see Ghilardi et al., 2010, table 1). Additionally, *P. mezzalirai* differs from these other fossil species in having more elliptical outline and in having a shallow suture. From the recent physid species (Simone, 2006: 181), *Physa mezzalirai* differs mainly by the wider body whorl, by shallower suture, by more spiral whorls, by narrower aperture, by sharper pointed spite, and by micro-sculpture in the body whorl (Fig. 8).

## Acknowledgments

We wish to thank Philippe Bouchet (Muséum National d'Histoire Naturelle, Paris, France) by the attention in informing us about the ICZN rules. By Tania Dutra, Editor of Gaea, for permission of publishing this additional and complementary paper.

## References

- GHILARDI, R.P.; CARBONARO, F.A. & SIMONE, L.R.L. 2010. *Physa mezzalirai*: um novo gastrópode da Formação Adamantina (Bacia Bauru), Cretáceo Superior, São Paulo, Brasil. Gaea 6(2): 63-68.
- MEZZALIRA, S. 1974. Contribuição ao conhecimento da estratigrafia e paleontologia do Arenito Bauru. Boletim do Instituto Geográfico e Geológico 51:1-163.
- SIMONE, L.R.L. 2006. Land and freshwater molluscs of Brazil. São Paulo, EGB. Fapesp, 390 pp.
- TAYLOR, D.W. 2003. Introduction to Physidae (Gastropoda: Hygrophila): biogeography, classification, morphology. Revista de Biología Tropical 51(suppl. 1):1-287.

Received: October, 2011. Accepted: November, 2011.